

Cunninghamia

A journal of plant ecology for eastern Australia

Date of Publication:
February 2018



ISSN 0727-9620 (print) • ISSN 2200-405X (Online)

Hydrogeomorphology, floristics, classification and conservation values of the little-known montane mires of the upper Cudgegong River catchment, Central Tablelands, New South Wales

Ian R. C. Baird^{1,2} and Doug Benson³

¹School of Health and Science, Western Sydney University, Locked Bag 1797, Penrith, NSW 2751, AUSTRALIA.

²Current address: 3 Waimea St, Katoomba, NSW 2780, AUSTRALIA. petalurids@gmail.com

³Honorary Research Associate, National Herbarium of New South Wales, Botanic Gardens & Domain Trust, Sydney 2000, AUSTRALIA. Doug.Benson@rbgsyd.nsw.gov.au

Abstract: Mires or peat swamps have a restricted distribution in Australia and are limited to areas where hydrological inputs exceed evapotranspiration. In NSW, mires are restricted to the coast, adjacent ranges or tablelands, and along the Great Dividing Range; most are listed as threatened ecological communities under State or Commonwealth legislation. Due primarily to the relatively high rainfall and suitable geology, the Blue Mountains region includes a number of such threatened mire ecological communities. Most of these mire types are largely included within the Greater Blue Mountains World Heritage Area, although there are notable exceptions, such as the endangered Newnes Plateau Shrub Swamps.

This paper reports on a little-known group of diverse, relatively isolated and largely unprotected mires, in a relatively low rainfall area in the upper Cudgegong River catchment, east of Rylstone in the NSW Central Tablelands, and of their floristic, hydrogeomorphic and typological relationship with other mires of the Blue Mountains. They can be broadly divided into montane bogs, montane fens and hanging swamps. Particular attention is focussed on the largest and most diverse one, Rollen Creek swamp, which contains all three types. It is hoped that highlighting this hitherto unrecognised group of high conservation-value mires will contribute to their improved conservation and encourage further research into mires of eastern NSW.

Key words: peat swamp, wetlands, groundwater-dependent ecosystems, *Eucalyptus camphora*, *Petalura gigantea*, National Heritage, Greater Blue Mountains World Heritage Area.

Cunninghamia (2018) 18: 001–021

doi:10.7751/cunninghamia.2018.18.001

Introduction

Wetlands encompass a range of vegetated ecosystems, including those referred to as mires, bogs, fens, swamps and marshes (see Gore, 1983; Mitsch & Gosselink, 2007; Mitsch et al., 2009). Wetlands are characterised by a diversity of hydrological regimes, and may be permanently, seasonally or intermittently inundated or saturated (DECCW, 2010). Even in permanent wetlands, water table depth can vary considerably within a particular wetland and between different wetland types. Such spatial heterogeneity, even within a permanent wetland complex, may result in considerable heterogeneity in substrates, vegetation associations and habitat, often across small spatial scales (e.g., Brown et al., 1982; Keith & Myerscough, 1993; Keith et al., 2006). For example, wetland complexes are often characterised by a complex intergrading of fens, bogs, swamps or marshes (Hájek et al., 2006; Kirkpatrick & Bridle, 1998; Rydin & Jeglum, 2013; Wheeler & Proctor, 2000; Yabe & Onimaru, 1997).

Hydrology and water balance (evapotranspiration compared to precipitation and other hydrological inputs) are the critical factors in determining the development of peaty or organic-rich wetland sediments. A basic requirement for peat formation is that plant biomass production (carbon production) exceeds decomposition (ecosystem respiration or carbon output). Consistently high water tables and a relatively anoxic environment generally provide the necessary conditions for peat accumulation in wetlands. Conditions of seasonal drying or widely fluctuating water tables, and/or negative water balance, result in oxidisation and bacterial decomposition of organic matter, and are not conducive to accumulation of peat (Gore, 1983; Rydin & Jeglum, 2013).

In the international context the term *mire* refers to peat-forming wetlands, and includes bogs and fens (Gore, 1983; Joosten et al., 2017; Rydin & Jeglum, 2013), although the terms may be applied somewhat differently in Australia (see Whinam & Hope, 2005). In the Australian context, bogs are typically low nutrient, acidic, dominated by sclerophyllous sedges and shrubs, and mosses, and may have large parts of the surface raised above the water table, while fens are usually more nutrient-rich, less acidic or alkaline, dominated by softer herbaceous and graminoid vegetation, and usually with a near surface water table (Keith, 2004; Whinam & Hope, 2005). Additionally, in Australia, the term *swamp*, as used in this paper, is a generic term which may refer to a wide range of wetland types, including peat-forming wetlands (such as bogs and fens), and wetlands with primarily mineral substrates with environmental conditions unsuitable for peat development, in contrast for example, to its more specific application in the USA (Cowardin et al., 1979) or Europe (Rydin & Jeglum, 2013). While they receive hydrological inputs from precipitation, surface flow and groundwater in varying proportions, most mires (peat swamps) in Australia, including the diverse mire communities across the Blue Mountains of NSW, are considered groundwater-dependent ecosystems (see NSW Government, 2002; Serov et al., 2012; Whinam & Hope, 2005).

Mire ecosystems in the Blue Mountains are captured within the Montane Bogs and Fens and Coastal Heath Swamps vegetation classes of NSW and the ACT of Keith (2004). These are represented by a number of mire vegetation types, including Blue Mountains Sedge Swamps (BMSS), Newnes Plateau Shrub Swamps (NPSS), Cocks River Swamps and Boyd Plateau Bogs (Benson & Keith, 1990; Keith & Benson, 1988). These are characterised by considerable spatial heterogeneity across a number of environmental gradients, within and between individual swamps and swamp types (e.g., Holland et al., 1992a; Holland et al., 1992b). Variation in vegetation across the hydrological gradient (from ephemeral to permanent saturation) is particularly evident within the upland mires developed on sandstone geology (e.g., Benson & Baird, 2012; Holland et al., 1992a). The term *montane*, as used in this paper, follows its usage in various vegetation publications in NSW (e.g., Bell et al., 2008; Hunter & Bell, 2007; Keith, 2004; NSWSC, 2004). In describing the Montane Bogs and Fens vegetation class, Keith (2004) indicated an elevation range of 600–1500 m, which includes the elevation range of the study area. Following Keith (2004), bogs and fens identified in this study are thus referred to as montane bogs and montane fens. In their classification of the native vegetation of southeast NSW, Tozer et al. (2010) attributed higher elevation mire vegetation in the southern half of the Blue Mountains (the northern extent of their study area does not extend beyond Lithgow) to the Tableland Bogs, Tableland Swamp Meadows (in the Montane Bogs and Fens vegetation class) and Blue Mountains-Shoalhaven Hanging Swamps (in the Coastal Heath Swamps vegetation class) types.

The upper Cudgegong River catchment (east of Rylstone) on the NSW Central Tablelands drains an area surrounded by the Great Dividing Range (at elevations over 1000 m) and forms part of the inland flowing Macquarie River catchment (Fig. 1). Part of the upper Cudgegong River catchment (in the Dunns Swamp area) is included in Wollemi National Park in the Greater Blue Mountains World Heritage Area (GBMWhA) (Fig. 2). No comprehensive vegetation or wetland mapping has been undertaken across this catchment, although previous reports have identified and briefly described a range of wetland vegetation types in the Dunns Swamp area (Bell, 1998a, b; Tame, 1997). During doctoral thesis fieldwork on the biology of the endangered mire-dwelling Giant Dragonfly, *Petalura gigantea*, Baird (2012) identified a number of previously undocumented and relatively isolated mires, with disjunct flora and fauna occurrences, in and adjoining the upper Cudgegong River catchment. Baird and Benson (2017) subsequently highlighted the value of one of these mire systems, Rollen Creek swamp, which occurs largely in Coricudgy State Forest, in support of a proposal to add Coricudgy State Forest to the National Heritage list; ultimately a candidate for potential addition to the GBMWhA (GBMWhAAC, 2015). The added inclusion of specific areas on the edges of the GBMWhA, including Coricudgy State Forest, would significantly enhance the currently established biodiversity

values of the GBMWSHA, and has been recommended (Benson & Smith, 2015).

The aim of this paper is to draw attention to the overlooked mires of the upper Cudgegong River catchment, with a particular focus on the Rollen Creek mire system, the largest and most diverse, to contextualise them in relation to other mires of the Blue Mountains, to contribute to improving their conservation, and to encourage further research on mires in southeastern Australia.

Upper Cudgegong River study area

Location and physiography

The study area is the upper Cudgegong River catchment between the Great Dividing Range and Rylstone, centred approximately on Dunns Swamp, Wollemi National Park (32° 50' 04" S, 150° 12' 21" E). The area is located in the NSW Central Tablelands, in the north-western part of the Blue Mountains and of the Sydney Basin bioregion (Thackway & Cresswell, 1995). The Great Dividing Range forms the watershed of the upper Cudgegong River, which drains inland to the Macquarie River (Fig. 1 & 2). Drainages to the north, east and south of the upper Cudgegong River watershed drain to the coast. Elevation decreases from east to west; from 1256 m on the basalt peak of Mt Coricudgy on the Great Dividing Range, westward to 571 m at Rylstone. Additional high elevation areas (above 1000 m) along the catchment watershed, and associated with the Great Dividing Range, include the remnant basalt-capped plateau of Nullo Mountain, and the isolated basalt-capped peaks of Mt Darcy and Mt Coorongooba.

The geology of the study area consists of underlying Permian sandstone, conglomerate, shale and siltstone; Triassic Narrabeen Sandstone, shale and conglomerate, forming characteristic 'pagoda country'; and Tertiary basalt-capped peaks and remnant plateaus. The Capertee, Collingwood, Inglewood, Lees Pinch, Munghorn Plateau, Nullo Mountain and Coricudgy soil landscapes dominate the study area (Kovac & Lawrie, 1991). The Inglewood (Yellow Earths), Lees Pinch (Shallow Soils) and Munghorn Plateau (Siliceous Sands) soil landscapes have developed primarily in association with the Triassic Narrabeen sandstone geology, while Nullo Mountain and Coricudgy soil landscapes (Krasnozems) have developed in association with the basalt-capped peaks and remnant plateaus, where *in situ* weathered Tertiary basalt and basalt colluviums overlie the Narrabeen sandstone. Capertee (Yellow Podzolics) and Collingwood (Red Podzolics) soil landscapes have developed in association with the Permian geology (Kovac & Lawrie, 1991). The peaty sediments of the mires in the area were not differentiated in the coarse-scale soil landscape mapping of Kovac and Lawrie (1991).

The study area includes the Dunns Swamp area of Wollemi National Park, a popular tourist site, and parts of Nullo Mountain and Coricudgy State Forests. With the exception of logging of better quality timber from the slopes of Mt Coricudgy, there is no evidence of recent logging activity in the immediate Coricudgy State Forest area which is mainly essentially undisturbed shrubby woodland with little timber value. Variably cleared and farmed freehold lands are mainly concentrated along the Cudgegong River valley and its tributaries, such as Coxs Creek, and in the Nullo Mountain area. The lower Cudgegong valley around Rylstone was explored by the botanist Allan Cunningham in the 1820s, and the area taken up for pastoral settlement soon after. Occupational licences in the upper Cudgegong in the Parish of Kelgoola were offered for sale in 1843 (*Sydney Morning Herald* 18/1/1843).

Climate

Rainfall across the study area is characterised by high inter-annual variability, with average annual rainfall decreasing with elevation and from east to west. Highest mean monthly rainfalls occur between November and March (late spring to early autumn), with lowest mean monthly rainfalls from July to September (winter to early spring). Mean annual rainfall at the Nullo Mountain AWS (1130 m elev.) was 955 mm/annum (1994–2017) (www.bom.gov.au). Mean maximum monthly temperature at Nullo Mountain was 16.7° C (9.2–24.0° C) and mean minimum monthly temperature was 8.0° C (2.5–13.6° C). Mean Annual rainfall at the property "Kelgoola" (747 m elev.) in the upper Cudgegong River valley near Rollen Creek swamp and Mt Coricudgy, was 809 mm (1963–2006), also with high inter-annual variability (486–1235 mm/annum). The higher elevation areas along the Divide, including Mt Coricudgy, Mt Coorongooba and Mt Darcy, can also be expected to have higher rainfall than that of the nearby upper Cudgegong valley at "Kelgoola" due to an orographic effect, and similar to that of Nullo Mountain. Mean annual rainfall at Rylstone (Ilford Rd AWS, 605 m elev.), at the western edge of the study area, is lower, with 655 mm/annum. Mean maximum monthly temperature at Rylstone was 22.6° C (14.4–30.7° C) and mean minimum monthly temperature was 8.2° C (1.2–15.9° C) (www.bom.gov.au).

Previous vegetation studies in the upper Cudgegong

Previous vegetation studies in the upper Cudgegong are mainly confined to the Dunns Swamp area (Bell, 1998a; Gellie & McRae, 1985; Tame, 1997). Gellie and McRae (1985) referred more broadly to the Cudgegong Swamps. Building on a brief report by Tame (1997), and with limited additional fieldwork, Bell (1998a) described Cudgegong River Swamp Grassland, Upper Cudgegong Alluvial Sedgeland, Upper Cudgegong Alluvial Reedland, Upper Cudgegong Alluvial Shrub-swamp and Upper Cudgegong Sphagnum Bog in the Dunns Swamp area of Wollemi National Park.




Fig. 1: Map of the Greater Blue Mountains World Heritage Area showing the location of the upper Cudgegong River catchment east of Rylstone (inset box).




Fig. 2. Map of the Upper Cudgegong River catchment study area and locations of surveyed mires. The site codes in this diagram are explained in Table 1.

Although Bell (1998a, b) did not identify the *Carex* fen vegetation along Never Never Creek and other similar sites, he noted the need for additional survey of wetland vegetation communities across the upper Cudgegong area to clarify their floristic relationships and distributions. Bell (1998a) noted that the Cudgegong River Swamp Grassland vegetation occurred only in waterlogged peaty alluviums and the Upper Cudgegong Alluvial Sedgeland occupied “grassy bogs”. Tame (1997) suggested the substrate of this vegetation type was probably a shallow sandy peat overlying impermeable clays. Bell (1998b) noted that Cudgegong Sphagnum Bogs only occurred in limited areas on heavy soils with poor drainage, along creek-lines or in small localised patches, but also referred to it occurring on peaty sands.

Bell (1998a) also noted the mention by Ford (1989) of an area of swamp near Mt Darcy (“along Rollen Creek”) with apparent similarities to his Cudgegong River Swamp Grassland. This is the first known reference to a swamp system along Rollen Creek, later described by Baird (2012) and Baird and Benson (2017).

Methods

Reconnaissance fieldwork (by IRCB) was carried out across the upper Cudgegong River catchment and adjoining areas in 2007-2009, searching for evidence of the endangered *Petalura gigantea* and to identify mires with potential breeding habitat for this obligate, groundwater-dependent, mire-dwelling dragonfly (Baird, 2012, 2014). Identification of upper Cudgegong mire sites has been based on a range of sources; aerial photograph interpretation (API, using Google Earth); previous vegetation reports from the Dunns Swamp area (Bell, 1998a, b; Tame, 1997); additional information provided by former NPWS ranger Chris Pavich (pers. comm.); data gathered by Baird (2012) (which included recording mires along 4 creek systems), and the results of a vegetation survey of the swamp system along Rollen Creek in February 2017 (Baird & Benson, 2017); and additional vegetation surveys across the study area in October 2017. Swamps were surveyed across Coxs, Ganguddy, Kings Swamp, Never Never and Rollen creeks (Fig. 2). Swamps on the remaining creek systems are on private property. Vegetation surveys consisted of walking transects targeted to cover as much local variation as possible and to record all readily identifiable plant species (and any evidence of threatened faunal species) across the range of identified floristic and hydrogeomorphic variability within the swamps surveyed. The presence of localised seepage areas or of an emergent water table provided further confirmation of the presence of suitable hydrological conditions for the development of organic-rich substrates. A sediment probe (1.8 m long × 8 mm diameter steel rod) was used to measure depth, and qualitatively assess characteristics, of organic-rich sediments in lower Rollen Creek swamp.

Results

Mires of the upper Cudgegong River catchment

Swamp vegetation in the upper Cudgegong River catchment was identified along the Cudgegong River and its tributary creeks - Coxs, Dairy Swamp, Ganguddy, Gavins, Kings Swamp, Mill, Never Never, Rollen, Sugarloaf and Swampy creeks (Fig. 2). Mire vegetation communities (with persistent high water tables and organic-rich or peaty sediments) were confirmed by survey along Coxs, Ganguddy, Never Never and Rollen creeks. Mire vegetation was identified along the main Cudgegong River valley immediately upstream and downstream of the confluence with Rollen Creek, but was not surveyed due to its location on private property. The presence of mire was not confirmed along the highly degraded Kings Swamp drainage. Details of mires surveyed are shown in Table 1. Occasional small isolated patches of seepage-fed hanging swamp on valley sides are also scattered across the upper study area on Narrabeen Sandstone, particularly in the Rollen Creek catchment. Additional small mire patches can be expected to occur elsewhere across the study area in locations that are difficult to access or identify on Google Earth imagery.

The mires of the upper Cudgegong River and its tributaries above Dunns Swamp, and along upper Coxs Creek, occur primarily in the Munghorn Plateau soil landscape on Triassic geology (see Kovac & Lawrie, 1991). Mire sediments included organic sands, sandy peats, sapric peats and fibrous peats, depending on the hydrological regime and topographic context. Some patches of swamp with sedgeland-heath vegetation, with floristic similarities to the wetter sedgeland-heath bogs, occur on heavier clayey soils (e.g., along lower Ganguddy Creek above Dunns Swamp) and cannot be considered mires. Depths of soft, saturated, organic-rich/peaty sediments throughout the main valley floor along lower Rollen Creek swamp (CM01) were generally >1.8 m, frequently with no major differences in density or texture observed across that depth. Dense sandy-gravel material was encountered, however, below 1.6 m in several locations. In one area near the confluence with a tributary near the Coricudgy State Forest boundary, some narrow bands of sandy-gravel material at different depths indicated historical deposition events, probably associated with post-fire erosion, upstream and/or upslope. Predictably, depth of organic-rich swamp sediment progressively decreased, and sand content increased, towards swamp margins on lower valley side slopes.

Considerable hydrogeomorphic and floristic diversity was observed in the mires across the study area, including montane bogs and montane fens, broadly referable to the Tableland Bogs and Tableland Swamp Meadows typologies, respectively, of Tozer et al. (2010), within the Montane Bogs and Fens vegetation class of Keith (2004), and hanging swamps on valley sides, with similarities to the Blue Mountains-Shoalhaven Hanging Swamps typology of Tozer et al. (2010), within the Coastal Heath Swamps vegetation class of Keith (2004). The rare, swamp-dwelling small tree,

Eucalyptus camphora subsp. *camphora*, is widely distributed across the upper Cudgegong catchment within each of these mire variants, along associated upstream drainage lines, and in some other swampy areas on heavier soils, including along lower Ganguddy Creek.

The main mire systems

Rollen Creek swamp, most of which occurs in Coricudgy State Forest, is one of only two extensive and relatively undisturbed (based on API, using Google Earth) mire systems in the upper Cudgegong catchment. The Rollen Creek swamp system appears to include the full range of hydrogeomorphic and floristic variation observed within the identified mires in the study area, including montane bogs (Tableland Bogs) and montane fens (Tableland Swamp Meadows) on valley floors, and hanging swamps on valley sides (Figures 3-11). Due to its outstanding diversity, size, condition and its representativeness of the range of mires in the study area, the Rollen Creek swamp system is described in detail below.

In addition to the extensive area of valley-floor sedgeland-heath bog (Tableland Bog) along Rollen Creek, several isolated and very small patches of variably degraded, seepage-fed sedgeland-heath bog (CM08-10) were also identified adjacent to, and intergrading with, the alluvial *Carex*-dominated fen (Tableland Swamp Meadow) (CM07) along upper Coxs Creek (Fig. 13). A narrow band of hanging swamp (CM04) of sedgeland-heath and *Sphagnum*, with conspicuous springs, also occurs just above Ganguddy Creek, near to and upstream of the Coricudgy Road bridge.

The other large mire system in relatively good condition is along Never Never Creek and its lower tributaries above Dunns Swamp (Kandos Weir impoundment) in Wollemi National Park. The Never Never Creek mire system is dominated by *Carex* fen (Tableland Swamp Meadow) along the main valley floor (CM03) with scattered *Leptospermum* shrubs (Fig. 12), although floristic variants occur in some marginal areas and in slightly higher gradient tributary swamp (CM02), in association with *Eucalyptus camphora*. The Rollen Creek and Never Never Creek mire systems are surrounded by high quality, dry sclerophyll eucalypt-dominated woodland on adjoining slopes.

The swamps along the main valley floors of the Cudgegong River (above and below Dunns Swamp) and its tributaries, including Coxs Creek (Fig. 13) are generally low gradient, alluvial swamp meadows. They are generally located on cleared freehold land, with a history of grazing, nutrient enrichment and weed invasion, and mostly range in condition from moderately- to highly-degraded. They range from *Carex*-dominated fens with organic-rich alluvium in wetter areas (Tableland Swamp Meadows) which may be variably inundated with shallow water, to seasonally wet tussock grasslands on largely mineral soils at the drier end of the hydrological gradient. Some of the swamp meadows include small pools along the main drainage line (possible chain-of-ponds system) with reedland (marsh) vegetation

of *Typha* sp., *Phragmites australis* and/or *Eleocharis sphecelata*. Along the main Cudgegong River valley above Dunns Swamp, around the confluence with Rollen Creek, areas of highly degraded Tableland Swamp Meadow sometimes also appear to have small remnant patches of sedgeland-heath bog (Tableland Bog) on their margins. Based on API (using Google Earth), additional areas of Tableland Swamp Meadow with possible Tableland Bog were identified along Gavins and Sugarloaf creeks, upper tributaries of the Cudgegong River; these swamps on private properties are mostly surrounded by cleared grazing land and were not visited.

A localised part of the upper Kings Swamp drainage includes some plants which occur in mires elsewhere in the area (e.g., *Callistemon citrinus*, *Eucalyptus camphora*, *Leptospermum obovatum*), with some groundwater seepage evident, but it is highly degraded and occurs on cleared grazing land. Regardless of its potential pre-disturbance state, it is not treated as mire here. A degraded and grazed narrow swamp with *Callistemon citrinus* and *Eucalyptus camphora* subsp. *camphora* also occurs along upper Mill Creek. This small swamp patch on otherwise cleared, private property was observed but not surveyed and it was not possible to determine its substrate or hydrological characteristics. It is also not treated as mire here.

Another large swamp system is in Jones Hole, on Jones Hole Creek (a headwater tributary of Coricudgy Creek), to the northwest of Mt Coricudgy (Fig. 2), in the coastward flowing Hunter River catchment, not the Cudgegong River. It is mostly in Wollemi National Park though partly in Coricudgy State Forest. The vegetation of this difficult-to-access valley-floor swamp system (32° 48' 16" S, 150° 20' 11" E, 720–850 m elev., ~3250 m length) is unknown, but it is reasonable to assume, due to its elevation and proximity to the higher rainfall area of adjacent Mt Coricudgy, that it is characterised by a high water table and variably peaty sediments. API (using Google Earth) suggests that it is predominantly a *Carex*-dominated fen, with areas of intergrading sedgeland-heath bog. The swamps in Jones Hole and along Never Never Creek have some history of cattle grazing prior to National Park gazettal (C. Pavich, pers. comm.). Conspicuous, abundant and structural plant species recorded for each of the identified and surveyed mires in the study area are included in Table 2. This is an incomplete list as some grasses and monocots were not identified, and some seasonal species may not have been evident at the time. Only two conspicuous bryophytes are listed, but the total bryophyte richness is likely to be high.

Other swamps

The study area includes a diversity of swamp vegetation types distributed across the hydrological gradient; much of the swamp area is characterised by only seasonally or intermittently wet sandy alluviums or heavier clayey soils along valley floors and cannot be considered mires. This includes seasonally wet tussock grasslands on largely mineral

soils at the drier end of the hydrological gradient, which generally occur in the lower elevation and lower rainfall parts of the catchment. Most of these seasonally wet tussock grasslands are surrounded by cleared grazing areas and occur along the Cudgegong River and its tributaries below Dunns Swamp, and along lower Coxs Creek; they have a history of grazing, nutrient enrichment and weed invasion; and are moderately- to highly-degraded.

Some shrubby swamps on sandy alluviums, including in the Dunns Swamp area and upper Ganguddy Creek, have a scrub of *Leptospermum polygalifolium* (often colonising disturbed sites), and are probably referable to the Upper Cudgegong Alluvial Shrub-swamps of Bell (1998a). Upper Cudgegong Alluvial Reedland (fringing vegetation around the Kandos Weir impoundment) and Upper Cudgegong Alluvial Shrub-swamp of Bell (1998a) in the Dunns Swamp area are not mires and were not considered potential habitat for

P. gigantea by Baird (2012). Other reedland (marsh) patches, however, occur in association with the *Carex* fens and can be considered part of the mire systems. Swamps in the lower parts of the Cudgegong River catchment (east of Rylstone) below Dunns Swamp and along lower Coxs Creek, occur in the Capertee soil landscape on Permian geology (Kovac & Lawrie, 1991).

Additional seasonally wet swamps occur to the north of the catchment watershed in headwaters of the northward flowing Growee River catchment (e.g., Spring Log Swamp on Spring Log Creek), but these are on primarily mineral sediments and are not mires. Growee Swamp (500 m elevation), which occurs further downstream on the Growee River to the north, was not visited, but it occurs on cleared grazing land in a lower rainfall area, and is also assumed to be on seasonally drying alluvial sediments or heavier clayey soils.

Table 1: Identified mires of the upper Cudgegong River catchment (arranged by decreasing elevation) showing - site code, location, coordinates, approximate elevation range (upper to lower), approximate length or area, mire type and condition. An unsurveyed Cudgegong River mire above Dunns Swamp has been included. Mire types - Tableland Bog (TB), Tableland Swamp Meadow (TSM), and Hanging Swamp (HS). The locations of these mires are indicated in Fig. 2.

Mire site	Location	Coordinates (Google Earth)	Elevation (m) (Google Earth)	Length or area	Mire type	Condition
CM01	Rollen Creek	32° 52' 47" S, 150° 17' 01" E	745-700	3 km	TB, HS, TSM	Good
CM07	Coxs Creek	32° 44' 27" S, 150° 09' 56" E	740-660	10 km	TSM	Poor-Good
CM08	Coxs Creek margins adjoining CM07	32° 44' 43" S, 150° 10' 18" E	724	20 × 20 m	TB	Poor
CM09	Coxs Creek margins adjoining CM07	32° 44' 37" S, 150° 10' 07" E	722	40 × 40 m	TB	Poor
CM10	Coxs Creek margins adjoining CM07	32° 44' 35" S, 150° 09' 48" E	718	150 × 40 m	TB	Poor-Moderate
CM04	Above Ganguddy Creek	32° 51' 22" S, 150° 12' 52" E	680-678	100 × 5 m	HS	Good
CM05	Ganguddy Creek below and adjoining CM04	32° 51' 22" S, 150° 12' 52" E	678	12 × 10 m	TB	Moderate -Good
CM02	Never Never Creek tributary	32° 49' 26" S, 150° 12' 49" E	677-665	200 m	TB/TSM	Good
CM06	Ganguddy Creek tributary at junction with Ganguddy Creek	32° 51' 07" S, 150° 12' 51" E	674	15 × 10 m	TB	Moderate
CM03	Never Never Creek	32° 49' 32" S, 150° 12' 46" E	672-656	1.7 km	TSM	Good
Not surveyed	Cudgegong River above Dunns Swamp, upstream and downstream of the confluence with Rollen Creek	32° 05' 55" S, 150° 17' 04" E (indicative location)	741-719		TSM, TB	Very poor

Table 2: Native plant species (including mosses) recorded for each of the mires surveyed in the upper Cudgong River catchment. Refer to Table 1 for details.

SPECIES	Family	CM01	CM02	CM03	CM04	CM05	CM06	CM07	CM08-10
<i>Asperula gunnii</i>	Rubiaceae	1							
<i>Baeckea utilis</i>	Myrtaceae	1							
<i>Baloskion fimbriatus</i>	Restionaceae		1						1
<i>Baloskion australe</i>	Restionaceae	1							
<i>Baumea</i> sp.	Cyperaceae	1	1		1		1	1	1
<i>Blechnum nudum</i>	Blechnaceae	1			1	1			
<i>Bulbine bulbosa</i>	Liliaceae	1							
<i>Callistemon citrinus</i>	Myrtaceae	1	1	1	1	1	1		
<i>Carex appressa</i>	Cyperaceae							1	
<i>Carex gaudichaudiana</i>	Cyperaceae	1	1	1	1	1	1	1	1
<i>Centella asiatica</i> (or <i>cordifolia</i>)	Apiaceae	1							1
<i>Centrolepis?</i>	Centrolepidaceae	1							
<i>Comesperma retusum</i>	Polygalaceae	1							
<i>Drosera binata</i>	Droseraceae	1			1				1
<i>Drosera peltata/auriculata</i>	Droseraceae				1				
<i>Drosera spathulata</i>	Droseraceae	1							
<i>Eleocharis sphacelata</i>	Cyperaceae			1				1	
<i>Empodisma minus</i>	Restionaceae	1			1				
<i>Epacris microphylla</i>	Ericaceae	1			1				
<i>Epacris paludosa</i>	Ericaceae	1				1	1		
<i>Eriocaulon scariosum</i>	Eriocaulaceae	1							
<i>Eucalyptus camphora</i> subsp. <i>camphora</i>	Myrtaceae	1	1	1	1	1	1		1
<i>Eucalyptus pauciflora</i>	Myrtaceae	1							
<i>Gahnia sieberiana</i>	Cyperaceae				1				
<i>Geranium neglectum</i>	Geraniaceae	1	1						1
<i>Gleichenia dicarpa</i>	Gleicheniaceae	1			1	1			
<i>Gonocarpus micrantha</i>	Haloragaceae	1							
<i>Goodenia</i> sp.	Goodeniaceae	1			1				
<i>Gratiola peruviana</i>	Scrophulariaceae						1		
<i>Gymnoschoenus sphaerocephalus</i>	Cyperaceae	1							
<i>Hakea microcarpa</i>	Proteaceae	1							1
<i>Hybanthus?</i>	Violaceae	1							
<i>Hydrocotyle</i> sp. (<i>sibthorpioides</i> ?)	Apiaceae	1					1		
<i>Hypericum</i> sp. (<i>gramineum</i> ?)	Hypericaceae	1							
<i>Isachne globosa</i>	Poaceae	1							
<i>Isotoma fluviatilis</i> ?= <i>Pratia surrepens</i>	Campanulaceae	1							
<i>Juncus</i> spp.	Juncaceae	1		1				1	1
<i>Lepidosperma limicola</i>	Cyperaceae	1							
<i>Leptospermum continentale</i>	Myrtaceae	1	1		1				1
<i>Leptospermum grandifolium</i>	Myrtaceae				1	1	1		
<i>Leptospermum obovatum</i>	Myrtaceae	1	1	1		1	1		1
<i>Lepyrodia</i> sp.	Restionaceae	1			1				
<i>Lythrum salicaria</i>	Lythraceae							1	
<i>Patersonia fragilis</i>	Iridaceae	1							
<i>Phragmites australis</i>	Poaceae		1	1				1	
<i>Polytrichum</i> sp. (<i>Dawsonia</i> sp.?)	Polytrichaceae					1			
<i>Pultenea divaricata</i>	Fabaceae	1							
<i>Pultenea</i> sp.	Fabaceae	1							
<i>Ranunculus</i> sp.	Ranunculaceae	1							
<i>Senecio</i> sp.	Asteraceae	1							
<i>Sphagnum cristatum</i>	Sphagnaceae	1			1	1	1		1
<i>Spiranthes australis</i>	Orchidaceae	1							
<i>Stylidium graminifolium</i>	Stylidiaceae	1							1
<i>Tetrarrhena juncea</i>	Poaceae	1							
<i>Typha</i> sp. (<i>orientalis</i> ?)	Typhaceae							1	
<i>Utricularia dichotoma</i>	Lentibulariaceae	1							1
<i>Viola caleyana</i>	Violaceae	1				1			
<i>Xyris ustulata</i>	Xyridaceae	1							
<i>Xyris gracilis</i>	Xyridaceae	1							

Rollen Creek mire system: location and physiography

Because of its size, range of mire variation and good condition, the mire system along Rollen Creek deserves particular attention. Apparently once known as Rotten Creek (G. Summers pers. comm.), Rollen Creek is an upper Cudgegong River tributary about 13 km east of Olinda. The upper swamp is located in Coricudgy State Forest, with the contiguous downstream section of good quality swamp on freehold land ("Inglewood") extending to where the Coricudgy Road crosses (32° 52' 18" S, 150° 16' 59"E). Though cattle have historically had some access to at least parts of the swamp upstream of the Coricudgy Road crossing, it is generally unsuitable for cattle, largely due to its high water table and soft deep peaty soils (G. Summers pers. comm.). A somewhat degraded area of sedgeland-heath bog remnant also occurs downstream of the Coricudgy Road crossing (between the road and cleared grazing land) towards the Cudgegong River junction, but has been impacted by grazing, fire and a previous timber mill with associated sawdust dump which was located below the crossing (G. Summers pers. comm.).

The Rollen Creek mire system (CM01) is narrow and elongate, oriented roughly NNW-SSE along the creek valley, about 3 km long (about 2.5 km in the State Forest) and varies in width from about 10 to 100 m, but is mostly <40 m wide. The catchment is Narrabeen Sandstone with the exception of the isolated peaks of Mt Coorongooba (~1060 m) and Mt Darcy (1079 m), with their residual basalt, Coricudgy soil landscapes. Approximately 20% of the slopes of Mt Coorongooba, and almost 50% of the slopes of Mt Darcy are in the Rollen Creek catchment. The area of the catchment is approximately 18 km², but with the complexity of the hydrogeology of these Narrabeen sandstones, aquifers may be collecting water from a larger area. There is a conspicuous seepage/spring at the head of the main creek and several small drainage lines enter the swamp system along its length; some also have areas of groundwater-fed peaty swamps with sedgeland-heath vegetation along their lower sections where they join Rollen Creek.

The head of Rollen Creek mire is about 2 km from the Divide (to the south); Mt Darcy and Mt Coorongooba are both within 3 km of the head of the mire. Elevation ranges from about 745 m at the source of the mire to 700 m at the road crossing, giving a low overall gradient of 1.5% (15 m per km) and within the range of Newnes Plateau Shrub Swamps (NPSS) (Benson & Baird, 2012). The downstream end lacks a nick-point waterfall, characteristic of NPSS, but grades into a *Carex gaudichaudiana*-dominated fen above the Coricudgy Road crossing, and transitions to cleared grazing land downstream of the Coricudgy Road crossing, with the exception of the patch of remnant sedgeland-heath bog below the Coricudgy Road. The extent of the fen area is likely to have been increased (with conversion from sedgeland-heath bog) by damming associated with the road

crossing. The mire system is surrounded by dry eucalypt woodland on poor sandy soils on adjoining slopes.

Rollen Creek mire system: vegetation structure and composition

The Rollen Creek mire system is dominated by extensive, valley-floor, sedgeland-heath bog, referable to the Tableland Bogs typology. In addition to the seepage/spring at the head of the main creek, there are small seepage/spring-fed mire patches adjacent to the main valley-floor mire in at least two other locations further downstream. There is also an extensive valley-side hanging swamp dominated by *Gymnoschoenus sphaerocephalus*.

The Tableland Bog along the main Rollen Creek valley floor includes areas of closed and open sedgeland, with a variable proportion of shrubs forming either sedgeland-heath or with a mallee or tall shrub canopy (Figures 3–10). The swamp is essentially treeless, though much of it is dominated by areas of mallee or multi-stemmed, shrubby *Eucalyptus camphora* subsp. *camphora* plants. Towards the head of the mire system, where the valley narrows, older *Eucalyptus camphora* occasionally reach 6–8 m high. In this area, this mallee/tall shrub canopy (sometimes low open woodland) generally forms dense scrub associated with *Callistemon citrinus* and *Leptospermum* shrubs up to 5 m high. Along the lower parts of the swamp system, however, *Eucalyptus camphora* plants are much smaller and often less than 2 m high. Plants with a range of sizes/ages indicate continual recruitment of these species. Parts of the upper swamp system with a canopy of *Eucalyptus camphora* and other large shrubs also have a wet meadow/forb-land ground-layer, which includes various herbs, *Carex gaudichaudiana*, *Sphagnum cristatum* and unidentified grasses (Fig. 3). A similar vegetation variant was also recorded along a small tributary (CM02) adjoining the *Carex* fen along Never Never Creek (CM03), along lower Ganguddy Creek near the Coricudgy Road bridge (CM05, CM06), and in association with small patches of sedgeland-heath bog (CM08–10) along margins of the *Carex* fen along upper Coxs Creek (CM07).

Throughout the sedgeland-heath dominated valley-floor bog there may be an open or sparse, small-medium height shrub layer, including *Baeckea utilis*, *Comesperma retusum*, *Epacris microphylla*, *Epacris paludosa*, *Hakea microcarpa*, *Leptospermum obovatum* and *Pultenea divaricata*. The groundcover is generally very dense, mostly >90% cover, predominantly of *Sphagnum* moss, sedges and other graminoids, smaller shrubs and herbs, including carnivorous species such as *Drosera binata* and *Utricularia dichotoma*. Herbs and ferns include *Centella asiatica*, *Geranium* sp., *Gleichenia dicarpa*, *Goodenia* sp., *Isachne globosa*, *Spiranthes australis* and *Viola caleyana*. Sedgeland species within the sedgeland-heath include *Baumea* sp., *Baloskion australe*, *Empodisma minus*, *Gymnoschoenus sphaerocephalus*, *Juncus* sp., *Lepyrodia*

sp., *Tetrarrhena juncea* and *Xyris ustulata*. There are patches with *Sphagnum* hummocks up to 1 m high and the soil surface is spongy on flat and steeper side slopes (Figures 3, 10). There may also be localised areas of *Sphagnum* bog in shadier, wetter and more fire-protected sites along tributary drainage lines.

Carex gaudichaudiana is scattered throughout the valley floor mire in localised patches of fen amongst sedgeland-heath bog in wetter low gradient areas along the drainage lines (Fig. 6), and as a discrete patch of *Carex*-dominated fen referable to Tableland Swamp Meadow at the downstream end near the Coricudgy Road crossing (Fig. 11).

In hanging swamps on side slopes open patches of sedgeland with few or no shrubs may have *Gymnoschoenus sphaerocephalus* and *Gleichenia dicarpa* dominant, with *Empodisma minus*, *Lepidosperma limicola* and *Xyris ustulata* less abundant (Figures 8–10). *Baeckea utilis* generally occurs as scattered individuals across these *Gymnoschoenus*-dominated hanging swamp areas, with occasional other shrubs present. Upper hanging swamp margins are relatively distinct with a band of *Gleichenia* along the edge of the dry eucalypt woodland; *Eucalyptus camphora* also occurs on swamp margins.

The main woodland dominants adjoining the swamps are *Eucalyptus dives*, *Eucalyptus radiata*, *Eucalyptus rossii* and *Eucalyptus dalrympleana*. In one place (32° 53' 24" S, 150° 17' 24" E), the geographically restricted *Eucalyptus corticosa* was recorded in woodland adjacent to the upper margin of the hanging swamp on the eastern side of the Rollen Creek mire. *Eucalyptus pauciflora* may occur on woodland margins sometimes edging into the swamp, as it does in the higher elevation mires of the Newnes and Boyd plateaus further south.

About 48 native species, including most of the conspicuous, abundant and structural species, were recorded in Rollen Creek mire (Table 2). This is an incomplete list as some grasses and monocots were not identified, and some seasonal species may not have been evident at the time. Appendix 1 includes the list of species recorded in Rollen Creek swamp with reference to species also recorded in NPSS (Benson & Baird, 2012; Benson & Keith, 1990) and Boyd Plateau Bogs (Keith & Benson, 1988; Kodela et al., 1996).

Rollen Creek mire system: hydrology

Given the relatively low average annual rainfall compared to the nearby basalt-capped mounts and the higher elevation parts of the Blue Mountains further south, and its high inter-annual variability at the property “Kelgoola”, the presence of the well-developed mire system along Rollen Creek indicates a strong groundwater influence, in addition to rainfall input. This is exemplified by the presence of springs and hanging swamps with groundwater seepage adjacent to the valley floor mire. The Narrabeen Sandstone of the upper Blue Mountains is also characterised by complex

groundwater hydrology and the presence of aquifers which support a diversity of groundwater-dependent swamp communities (e.g., Baird, 2012; DLWC, 1999a, b; Marshall, 2005). The expected higher rainfall in the higher elevation areas above 1000 m, around the catchment watershed in the headwaters of the upper Cudgegong River catchment, is likely to contribute significantly to the presence of the groundwater-dependent mires through both direct runoff, and infiltration into the groundwater system in the relatively permeable geology.

The characteristic swamp vegetation and extensive occurrence of peaty/organic-rich mire sediments indicates a permanently high water table in Rollen Creek mire. Peaty sediments include fibrous peats and more highly decomposed sapric peats, which require such hydrological conditions to develop. Depth and characteristics of the peaty/organic-rich sediments can be expected to vary considerably across the swamp system, as occurs in NPSS for example, but the depths recorded in the downstream part of Rollen Creek swamp (>1.8 m) typically exceeded depths recorded in NPSS by Benson and Baird (2012). They are, however, similar to depths recorded in the somewhat similar, low gradient montane bog and fen complex at the head of Long Swamp (part of the Coxs River Swamps of Benson & Keith, 1990) in Ben Bullen State Forest (Baird, 2014; IRCB, unpubl. obs.; Martin, 2017).

The hanging swamps on the valley side, dominated by *Gymnoschoenus sphaerocephalus*, and contiguous with the main valley floor mire (Figures 8-10), are similar to those which occur elsewhere in the Blue Mountains on sandstone geology. These hanging swamps are typically associated with the presence of a low-permeability aquitard (e.g., claystone layer). They develop where groundwater is redirected downslope following the slope of the aquitard to emerge on the valley side as seepage. These hanging swamps along Rollen Creek include groundwater seeps and springs with associated groundwater-dependent species such as burrowing crayfish (*Euastacus* sp.) (Fig. 10).

The predominance of *Sphagnum* in the Rollen Creek mire system is indicative of persistent or permanent wetness. The hummocks are slow growing and the large hummocks were associated with large shrubs with a cluster of basal stems, or large tussocks of sedgeland vegetation, which appear to provide support for the hummock growth and provide moderate levels of shade. Larger *Sphagnum* hummocks in Boyd Plateau Bogs are similarly associated with multi-stemmed shrubs and sedgeland tussocks (IRCB unpubl. obs.). The *Sphagnum* is vulnerable to fire during drought, when the hummocks may dry out (see Hope et al., 2009; Whinam, 1995; Whinam & Chilcott, 2002; Whinam et al., 1989).



Fig. 3: Dense *Sphagnum* cover and hummocks in *Carex* and grass-dominated area in the upper section of Rollen Creek swamp, with *Eucalyptus camphora* subsp. *camphora*. Photo by Ian Baird



Fig. 4: Lignotuberous *Eucalyptus camphora* subsp. *camphora* resprouting after fire along a drainage line through a *Gymnoschoenus*-dominated patch of swamp in the upper Rollen Creek. Photo by Ian Baird



Fig. 5: Dense montane sedgeland-heath bog along the Rollen Creek valley floor. Photo by Ian Baird



Fig. 6: A patch of *Carex gaudichaudiana* fen amongst sedgeland-heath in a low gradient section of Rollen Creek swamp. Note the emergent water table (bottom centre) and *Eucalyptus camphora* subsp. *camphora* in front of adjoining woodland at rear. Photo by Ian Baird



Fig. 7: Valley floor bog with a patch of *Empodisma minus* and *Tetrarrhena juncea*-dominated ground-layer, with low *Eucalyptus camphora* subsp. *camphora* (rear and right). Photo by Ian Baird



Fig. 8: *Gymnoschoenus*-dominated hanging swamp in the upper section of Rollen Creek. The fringing low *Eucalyptus camphora* subsp. *camphora* is in front of adjoining upslope woodland at rear. Photo by Ian Baird




Fig. 9: Resprouting *Gymnoschoenus sphaerocephalus* tussocks post-fire in a hanging swamp in the upper Rollen Creek swamp. Note the large size (and age) of the bases of tussocks, fire-killed shrubs and fringing resprouting *Eucalyptus camphora* subsp. *camphora* (rear). Photo by Ian Baird



Fig. 10: Seepage in a *Gymnoschoenus*-dominated section of a hanging swamp in the upper Rollen Creek with *Sphagnum*, *Gleichenia dicarpa* and scattered shrubs of *Baeckea utilis* evident. A *Euastacus* sp. crayfish burrow is visible in the pool. Photo by Ian Baird



Fig. 11: *Carex gaudichaudiana* –dominated montane fen (centre left) in a low gradient area of the downstream end of the Rollen Creek swamp, with fringing sedgeland-heath bog. Photo by Ian Baird



Fig. 12: Valley-floor *Carex* fen along Never Never Creek above Dunns Swamp, with scattered *Eucalyptus camphora* subsp. *camphora* and *Leptospermum* shrubs. Photo by Ian Baird



Fig. 13: Valley-floor *Carex* fen with flowering *Lythrum salicaria* along Cocks Creek (looking upstream) showing cleared grazing land on private property on right-hand (southern) side and good quality bushland on the left-hand side. Small patches of fringing sedgeland-heath bog, with some *Eucalyptus camphora*, occur in embayments along fen margins at right. Photo by Ian Baird

Discussion

Comparison of montane mires of the upper Cudgegong River catchment with other montane mires of the Blue Mountains

We found a range of mires with considerable floristic and hydrogeomorphic diversity across the upper Cudgegong River catchment. These mires can be broadly divided into montane bogs, montane fens and hanging swamps; all groundwater dependant ecosystems typically found in high rainfall areas. Keith (2004) indicated a rainfall range of 800–1500 mm/year for the Montane Bogs and Fens vegetation class and 1000–1500 mm/year for the Coastal Heath Swamps vegetation class. The Cudgegong mires are at the extreme lower limits of these ranges. The upper

Cudgegong montane bogs (Tableland Bogs) and montane fens (Tableland Swamp Meadows) occur in a lower rainfall area, and at relatively lower elevation, compared to similar Blue Mountains mires further south. Mires in Ben Bullen State Forest and on the Newnes Plateau are above 900 m (but most above 1000 m), compared to those at 660–745 m in the upper Cudgegong (Table 1). Mean annual rainfall for these central Blue Mountain mires generally exceeds 1000 mm, and on the Boyd Plateau and nearby areas (above 1100 m elevation) exceeds 1100 mm.

The hanging swamps of the upper Cudgegong River catchment, particularly along upper Rollen Creek, also occur in a lower rainfall area than similar hanging swamps on sandstone in the central Blue Mountains (BMSS), particularly those on the eastern side of the Blue Mountains at similar elevation, where mean annual rainfall also exceeds 1100 mm, compared to mean annual rainfall of 809 mm at “Kelgoola”, near Rollen Creek mire. In upper parts of the Blue Mountains, where the best developed hanging swamps on sandstone occur, mean annual rainfall may exceed 1300 mm. In lower elevation areas with considerably lower rainfall and higher evapotranspiration in the lower reaches of the upper Cudgegong River catchment east of Rylstone (mean annual rainfall <750 mm), to the north of the upper Cudgegong River catchment in the Growee River catchment, and to the south in the Capertee River catchment, suitable conditions for mire development are evidentially absent.

Floristically, the Rollen Creek mire is similar to NPSS with about 65% of the 48 recorded native species in common, particularly shrub species (see Appendix 1), though the predominance of *Eucalyptus camphora*, restricted mainly to these swamps, is a conspicuous point of difference. *Eucalyptus camphora* subsp. *camphora* is geographically restricted to Nullo Mountain and the upper Cudgegong River catchment, with a localised disjunct population in swamps in the Megalong Creek valley near Katoomba. It is also of note in being restricted mainly to swamp habitat; indeed there are very few *Eucalyptus* species that are found in wet or poorly drained sites and *Eucalyptus camphora* appears to be confined to this habitat. The only similar swamp-inhabiting eucalypt in the Sydney Basin bioregion (there are almost 100 eucalypt species in the the GBMWA; Benson & Smith, 2015) is the closely related *Eucalyptus aquatica*, which is found in analogous montane mires in the Southern Highlands (Penrose State Forest) (Shepherd & Keyzer, 2014).

The abundance of *Callistemon citrinus* along Rollen Creek is another conspicuous difference. This species does not occur in NPSS or Boyd Plateau Bogs, but is the only *Callistemon* found in BMSS (and some transitional to NPSS). Interestingly, *Callistemon pityoides*, the typical montane bog *Callistemon* (found in Boyd Plateau Bogs, some NPSS, and montane bogs elsewhere), was not recorded anywhere in the upper Cudgegong swamps, nor does it occur in BMSS. Other conspicuous NPSS species not so far recorded are *Celmisia* sp. (aff. *longifolia*), *Boronia deanei* subsp. *deanei* and *Grevillea acanthifolia* subsp. *acanthifolia*. *Celmisia* and *Boronia deanei* also occur in Boyd Plateau Bogs. These species may be more drought-sensitive and given the much less extensive

swamp areas on the upper Cudgegong, in comparison with the main Blue Mountains area further south, could have been extirpated by past periods of extensive drought, assuming they have been present in the past.

Although *Leptospermum grandifolium* was recorded in the small patch of hanging swamp above Ganguddy Creek (CM04) and a nearby small tributary bog (near and downstream of the bridge on the Coricudgy Road) (CM06), the apparent lack of the species in Rollen Creek swamp and other surveyed sites is noteworthy, considering that it is a common shrub in NPSS, BMSS and Boyd Plateau Bogs (interestingly, single plants of *Epacris paludosa*, which was otherwise only recorded from Rollen Creek swamp, were also recorded in CM05 and the nearby CM06). There is also a general lack of dominance by *Leptospermum* species, compared to their abundance in many NPSS. The only species recorded in Rollen Creek swamp was *Leptospermum obovatum*, which is common in Rollen Creek swamp, occurs in some NPSS, and is common in Boyd Plateau Bogs. *Leptospermum continentalis* also occurs on heavier soils on drier margins of swamps. *Baeckea utilis*, the only *Baeckea* species present, and widely distributed in Rollen Creek swamp, is a typical montane bog and wet heath species which occurs in and around Boyd Plateau Bogs and some higher elevation Newnes Plateau swamps, often on heavier soils, but does not occur in BMSS. *Baeckea utilis* is typically replaced by *Baeckea linifolia* in most NPSS and in the higher elevation Coastal Heath Swamps of the region such as BMSS.

Most other abundant species are similar to those occurring in NPSS, including *Lepidosperma limicola*, *Empodisma minus*, *Epacris paludosa* and *Leptospermum obovatum*. Although *Gymnoschoenus sphaerocephalus* occurs in small patches within the main valley floor bog and in upstream areas where the valley narrows, it is mostly restricted to the hanging swamps and seepage/spring-fed mire patches adjacent to the main valley floor mire. Rollen Creek was the only mire in the upper Cudgegong catchment where *Gymnoschoenus* was recorded, although it has been recorded just east of Mt Coricudgy, in a previously undocumented small hanging swamp (990 m elev.) where *Petalura gigantea* was recorded (Baird, 2012: Appendix 1, site PMC01). *Gymnoschoenus* is widely distributed in mires on sandstone elsewhere in the Blue Mountains (in NPSS and BMSS) and the Southern Highlands, and montane bogs of the Northern and Southern Tablelands (mostly on granite), but it appears to be absent from the granite-based Boyd Plateau Bogs. The only occurrence in Tableland Bog on granite in the Blue Mountains region which is known to one of the authors (IRCB) is in a small montane bog south of the Kowmung River near Trailers Mountain, although it may also occur in several small unsurveyed bog patches nearby. Floristically about 58% of the 48 recorded native species in Rollen Creek swamp are shared with the Boyd Plateau Bogs, including herbaceous species such as *Centella asiatica*, *Geranium* sp., *Hypericum* sp., and *Viola caleyana* (Appendix 1).

Areas of hanging swamp dominated by *Gymnoschoenus sphaerocephalus* (Figures 8–10) are not referable to any

of the swamp vegetation types in the upper Cudgegong area identified by Bell (1998a). With the exception of the occurrence of *Eucalyptus camphora* and *Baeckea utilis*, these hanging swamps are more similar floristically and hydrogeomorphically to those occurring elsewhere on Narrabeen Sandstone in the Blue Mountains, such as the BMSS of Keith and Benson (1988).

The presence of extensive areas of *Sphagnum cristatum* along Rollen Creek, often forming large hummocks within the sedges and shrubs, was noteworthy. In comparison with Rollen Creek swamp, *Sphagnum* is either very restricted or absent from NPSS and BMSS (but note the extensive and unusual *Sphagnum* cover in the fen/marsh in Goochs Crater in the upper Wollangambe River catchment and in a BMSS in McCrae's Paddock in Katoomba). The much more extensive *Sphagnum* cover in Rollen Creek swamp (and elsewhere in mires and other parts of the upper Cudgegong catchment) is surprising considering the lower rainfall recorded at nearby "Kelgoola", compared to NPSS and other central Blue Mountains mires. Its persistence is likely to be the result, at least in part, of a sustained groundwater influence and the nearby presence of higher elevation peaks upstream around the catchment watershed contributing additional rainfall inputs as a result of an orographic effect. According to Chris Pavich (pers. comm.), *Sphagnum* was more abundant in the upper Cudgegong before the drought of the 1930-40s, with a subsequent increase in fire and grazing probably contributing to its further disappearance (C. Pavich pers. comm.). Rollen Creek swamp, however, has not been subject to heavy grazing or frequent anthropogenic fire (G. Summers pers. comm.), which may have helped prevent loss of *Sphagnum*. *Sphagnum* is generally much more abundant in montane bogs on granite, such as those of the Boyd Plateau, and nearby areas on metasedimentary geology (e.g., near Mt Werong and in Jenolan State Forest), than in the swamps developed on sandstone in the Blue Mountains, such as the NPSS and BMSS (also see Downing et al., 2007; Whinam & Chilcott, 2002). Differences in fire history and climatic factors between these areas are likely to be contributing factors.

While there are small patches of *Sphagnum* bog in the upper Cudgegong which are broadly consistent with Bell (1998a)'s Cudgegong *Sphagnum* Bogs, these are treated here as part of a variable upper Cudgegong montane bog type. *Sphagnum* also occurs along various drainage lines (often on heavy soils) and in localised patches within the other swamp types (including areas transitional between *Carex* fen and sedgeland-heath bog) where suitable conditions occur. The Cudgegong River Swamp Grassland and Upper Cudgegong Alluvial Sedgeland of Bell (1998a) are also treated as part of the montane fens identified in this study. As a result of more extensive survey across the study area, and based on their broadly similar floristics and potential organic-rich substrates, as described by Bell (1998a) in the Dunns Swamp area, Baird (2012) noted that there is considerable gradation between these swamp vegetation types.

In summary, the Rollen Creek swamp includes extensive valley floor mire of intergrading bog and fen, with some

very small valley-side seepage/spring areas, and extensive valley-side hanging swamp dominated by *Gymnoschoenus*. The valley-side seeps, springs and hanging swamps are most similar to other such groundwater-dependent mire expressions developed on Narrabeen Sandstone in the Blue Mountains, such as the NPSS and BMSS. While these seeps and hanging swamps are hydrogeomorphically similar to similar mires developed on sandstone geology elsewhere in the Blue Mountains, they are characterised by a somewhat distinctive floristic assemblage. BMSS form part of the Coastal Heath Swamp vegetation class (NSWSC, 2007), although most are above the nominal 600 m upper elevation range indicated by Keith (2004) for the Coastal Heath Swamps. NPSS are considered transitional between the Coastal Heath Swamps and Montane Bogs and Fens vegetation classes (NSWSC, 2005b), with that transition occurring between Bell and Clarence. The valley floor mire along Rollen Creek, however, with areas of bog and fen, shows greater affinity, respectively, to the Tableland Bogs and Tableland Swamp Meadows of Tozer et al. (2010), within the Montane Bogs and Fens vegetation class. The valley-floor bogs along Rollen Creek and elsewhere in the upper Cudgong have floristic and hydrogeomorphic similarities with montane bogs of the Southern, Central and Northern Tablelands (e.g., Hunter & Bell, 2007; Tozer et al., 2010). The *Carex* fens along Coss Creek and Never Never Creek, at the downstream end of Rollen Creek swamp, and similar areas across the upper Cudgong River catchment, have affinities with the *Carex* fen vegetation of Northern NSW (Hunter, 2013; Hunter & Bell, 2009), the *Carex-Poa* fen vegetation of Long Swamp in Ben Bullen State Forest and nearby areas (part of the Coss River Swamps of Benson & Keith, 1990), the swampy meadows of the Central Tablelands (see Mactaggart et al., 2008; Mactaggart, 2008) and the Mountain Hollow Grassy Fens (DEC, 2006). This complexity highlights the value of this and other geographically isolated mires of the upper Cudgong River catchment.

Conservation value of Rollen Creek swamp and other remaining mires of the upper Cudgong River catchment

Mires are geographically restricted ecosystems in Australia and their extent and health have been considerably reduced and degraded since European settlement through urban and transport infrastructure development, agriculture, drainage, grazing, and more recently through mining impacts, particularly the impacts of subsidence from longwall coal mining, a Key Threatening Process in NSW (CoA, 2005, 2010; NSWSC, 2005a). Climate change and fire are also recognised as significant threats to these ecosystems (e.g., Baird & Burgin, 2016; CoA, 2010; Keith et al., 2014; Pemberton, 2005).

In the Sydney Basin bioregion, Temperate Highland Peat Swamps on Sandstone (THPSS) is listed as an Endangered Ecological Community (EEC) under the Commonwealth *Environment Protection and Biodiversity Conservation Act 1999* (CoA, 2005). THPSS also includes NPSS and BMSS. Under the NSW *Biodiversity Conservation Act 2016* (which replaced the NSW *Threatened Species Conservation Act*

1995) NPSS is also listed as an EEC (NSWSC, 2005b), and Blue Mountains Swamps (including the Blue Mountains Sedge Swamps of Keith and Benson (1988)) is listed as a Vulnerable Ecological Community (NSWSC, 2007). Although not specifically included in the description of THPSS, the authors consider that some of the mires of the upper Cudgong, particularly the Rollen Creek valley floor mire and hanging swamps, are clearly referable to the THPSS EEC, and have been simply overlooked because of lack of documentation. For a similar reason, none of the valley-floor bogs and fens in the upper Cudgong are specifically identified in the Montane Peatlands and Swamps of the New England Tableland, NSW North Coast, Sydney Basin, South East Corner, South Eastern Highlands and Australian Alps bioregions EEC determination (NSWSC, 2004). In circumscribing the Montane Peatlands and Swamps EEC, the NSWSC (2004) referred to swamps above 400–600 m elevation. The montane bogs and fens of the upper Cudgong are also referable to this EEC and are clearly an important part of the complex of endangered montane mire communities distributed across the tablelands and adjacent ranges of NSW.

Habitat for rare species-flora

The mires of the upper Cudgong River catchment are isolated from other montane mires. The nearest montane mires to the north occur at high elevation at Barrington Tops (Mort, 1983), although somewhat drier and more floristically impoverished wet heath vegetation, with some floristic affinity to the wet heath in the upper Cudgong mires, occurs on heavier soils on Coolah Tops (Binns, 1997). The nearest to the south occur on the Newnes Plateau and the adjacent Ben Bullen State Forest.

This isolation contributes to the significance of some rare and restricted plant species in the mires and adjacent woodlands of the upper Cudgong. For example, the distribution of the mallee *Eucalyptus camphora* subsp. *camphora*, illustrates the disjunct biogeography of many montane species restricted to specialised habitats. This taxon is geographically restricted to higher elevation mires and swampy drainage lines along this part of the Divide, with populations at Nullo Mountain and in the upper Cudgong River catchment (Rollen Creek swamp in particular has a large population), and a highly localised and disjunct population 90 km further south in the Megalong Creek Valley near Katoomba, where it occurs in swamp patches with different floristics on heavier clayey soil and colluviums on Permian geology (600 m elev.). This locality in the Megalong Creek valley is also noteworthy for the occurrence of the rare and locally endemic *Callistemon megalongensis* (Craven, 2009; Udovicic & Spencer, 2012) and *Callistemon purpurascens* (Douglas & Wilson, 2015).

Another rare eucalypt, *Eucalyptus corticosa*, a locally endemic tree species restricted to the upper Cudgong valley east of Rylstone, occurs in eucalypt woodland on shallow infertile soils on sandstone ridges and was recorded on the outer edge of the *Gymnoschoenus*-dominated hanging swamp along Rollen Creek. *Veronica blakelyi*, a small shrub restricted to the western Blue Mountains, near Clarence,

near Mt Horrible, on Nullo Mountain and in the Coricudgy Range, is listed as Vulnerable under the NSW *Biodiversity Conservation Act 2016* and was recorded in woodland adjoining Rollen Creek swamp. Additional rare plant species are likely to occur in the woodlands adjoining these swamps, including in Coricudgy State Forest.

Habitat for rare species-fauna

The mires of the upper Cudgegong, particularly Rollen Creek, may also provide habitat for specialist mire fauna. Baird (2012) considered that the upper Cudgegong mires, particularly the valley-floor bogs and *Gymnoschoenus*-dominated hanging swamps, provide potential habitat for the endangered dragonfly, *Petalura gigantea*. The record of a single *Petalura* in a small hanging swamp (one of several such previously undocumented swamps in close proximity) east of Mt Coricudgy in Wollemi National Park, a considerable distance from the nearest known populations on the Newnes Plateau (Baird, 2012), strongly suggests that a population occurs somewhere within the complex of mires in the upper Cudgegong. Rollen Creek swamp is the most likely habitat, although surveys on three occasions have not recorded the species.

Rollen Creek swamp also appears to provide suitable habitat for the endangered Blue Mountains Swamp Skink, *Eulamprus leuraensis*, which is only known from mid-upper elevation mires in the central Blue Mountains (BMSS) and the Newnes Plateau (NPSS). This groundwater-dependent species has not been recorded further north than the Newnes Plateau (Gorissen, 2016; LeBreton, 1996). Evidence of foraging and tunnelling activity of Swamp Rats, *Rattus lutreolus*, a common species of BMSS and NPSS, was also observed during fieldwork. Additional obligate mire-dwelling fauna are likely to occur here (e.g., skinks and invertebrate stygofauna), but more detailed surveys will be required to identify them.

A burrowing spiny crayfish, *Euastacus australasiensis*, is widely distributed in Blue Mountains mires (including hanging swamps), and *Euastacus* burrows and partial remains were observed by the authors in the Rollen Creek mire system (including hanging swamps) and (by IRCB) in the small hanging swamp east of Mt Coricudgy where *Petalura gigantea* was recorded. While these are likely to be *Euastacus australasiensis* (see McCormack, 2012), the possibility exists that an unidentified swamp-dwelling taxon is involved. In the forested headwaters of the Cudgegong River in Coricudgy State Forest, below Mt Coricudgy and upstream of Rollen Creek, a large, stream-dwelling, burrowing spiny crayfish species, *Euastacus vesper*, closely related to *Euastacus spinifer* of eastern drainages, has recently been described (McCormack & Ahyong, 2017). The discovery of this apparently highly-localised species provides further evidence of the high conservation value of Coricudgy State Forest and of the opportunities for further research in this area.

Value of mires for palaeo-ecological, evolutionary and climate studies

Previous palaeoecological studies of Blue Mountains mires, based upon pollen and charcoal analysis of radiometrically-dated sediment cores, have greatly increased our understanding of the developmental history of these mires and their past climates and vegetation, particularly since the Last Glacial Maximum (~21 000 years BP). While the oldest sediment core ages suggest some mires may have commenced development around 13 000 years BP, sediment cores from other swamps have maximum ages within the Holocene (e.g., Black et al., 2008; Chalson & Martin, 2009; Fryirs et al., 2014; Martin, 2017). The mires of the upper Cudgegong River catchment can be expected to be a similarly rich source of knowledge related to species distributions, the developmental history of these mires and of climatic change since the Last Glacial Maximum, and expand our existing understanding of vegetation change across southeastern Australia and the Sydney Basin during this period.

The biogeography of the mallee *Eucalyptus camphora* subsp. *camphora*, geographically restricted to localised high elevation patches of suitable habitat along the Divide, is similar to that of many montane species with disjunct populations restricted to specialised habitats. Such disjunct distributions are likely to reflect, or be the result, of past climatic fluctuations to some extent, either during the Last Glacial Maximum or as a result of previous glacial/interglacial cycles. For *Eucalyptus camphora*, two other subspecies have also been recognised; subsp. *relicta*, found further north at Guyra and in Queensland, and subsp. *humeana*, occurring from Wee Jasper south into Victoria. The distribution of these taxa, with their greater morphological and geographic variation, presumably indicates older geographic separation than that within subsp. *camphora*, and provides evidence of past climate-related divergence. Other montane species restricted to specialised habitats, such as mires, have similar biogeographic patterns; understanding these patterns is of considerable scientific interest, particularly in the context of a rapidly changing climate.

Conclusion

The upper Cudgegong River catchment includes a complex of endangered peat swamp or mire types which are geographically disjunct from their nearest neighbours and characterised by some distinctive floristic assemblages. These include areas of montane bog, montane fen and hanging swamp. The presence of rare species such as the mallee *Eucalyptus camphora* subsp. *camphora*, and the potential for endangered fauna such as *Petalura gigantea* to be present, further highlights their value. In the context of the relatively low rainfall where these groundwater-dependent mires occur (and the upper Cudgegong is at their climatic limits), there is a surprising hydrogeomorphic and floristic diversity across these different mire types. With their relatively low rainfall, these mires may be particularly vulnerable to climate change.

Historically there has been considerable loss and degradation of the mires in the upper Cudgong River catchment through land clearing and agriculture, beginning in the 1840s, and evidence of this damage provides a strong imperative to protect those examples that have survived. Conservation of these mires and their associated flora and fauna will benefit from further survey, mapping and biodiversity census. In addition to improved management of identified mires on private lands, including the possible use of biodiversity conservation covenants and incentives, improved recognition of the inherent values of these mires will be fundamental to their long-term conservation. Rollen Creek swamp is unique in the area, in terms of its size, floristic and hydrogeomorphic diversity, and good condition, and its conservation must be a priority. National Heritage listing of Coricudgy State Forest would provide a substantial foundation to highlight the values of this mire system and its surrounding woodland landscape, including the biodiversity associated with the significant, higher rainfall, basalt-capped island peaks adjoining the GBMWA. Such listing is a prerequisite for nomination of this and other high biodiversity areas for addition to the GBMWA and is recommended.

Acknowledgements

Gay Summers, owner of “Inglewood”, generously provided access to Rollen Creek swamp on various occasions and shared her local knowledge. Chris Pavich (NPWS) shared his extensive local knowledge and provided assistance with property access for IRCB during earlier fieldwork. Ahamad Sherieff (NSW Office of Environment and Heritage) kindly prepared the line map of the study area. Huw Evans (Local Land Services) provided assistance during an earlier fieldtrip. Rob McCormack reviewed the text on spiny crayfish and Scott Mooney and an anonymous reviewer are thanked for their useful comments which have improved the paper. Initial fieldwork by IRCB was supported by a Western Sydney University Post-graduate Research Award.

References

- Baird, I.R.C. (2012). The wetland habitats, biogeography and population dynamics of *Petalura gigantea* (Odonata: Petaluridae) in the Blue Mountains of New South Wales. PhD thesis, Western Sydney University, Australia. Available from <http://handle.uws.edu.au:8081/1959.7/509925>.
- Baird, I.R.C. (2014). Larval burrow morphology and groundwater dependence in a mire-dwelling dragonfly, *Petalura gigantea* (Odonata: Petaluridae). *International Journal of Odonatology*, 17, 101-121. doi:10.1080/13887890.2014.932312
- Baird, I.R.C. & Benson, D. (2017). Survey and estimation of biodiversity values in relation to National Heritage listing for Rollen Creek swamp, Coricudgy State Forest, Central Tablelands, NSW. Unpublished report. Available from <https://doi.org/10.13140/RG.2.2.24271.48800>.
- Baird, I.R.C. & Burgin, S. (2016). Conservation of a groundwater-dependent mire-dwelling dragonfly: implications of multiple threatening processes. *Journal of Insect Conservation*, 20, 165-178. doi:10.1007/s10841-016-9852-3
- Bell, D.M., Hunter, J.T. & Haworth, R.J. (2008). Montane lakes (lagoons) of the New England Tablelands Bioregion. *Cunninghamia*, 10, 475-492.
- Bell, S.A.J. (1998a). Wollemi National Park Vegetation Survey: A Fire Management Document. Community Profiles, Vol. 2. Unpublished report to NSW National Parks and Wildlife Service, Upper Hunter District.
- Bell, S.A.J. (1998b). Wollemi National Park Vegetation Survey: A Fire Management Document. Main Document, Vol. 1. Unpublished report to NSW National Parks and Wildlife Service, Upper Hunter District.
- Benson, D. & Baird, I.R.C. (2012). Vegetation, fauna and groundwater interrelations in low nutrient temperate montane peat swamps in the upper Blue Mountains, New South Wales. *Cunninghamia*, 12, 267-307. doi:10.7751/cunninghamia.2012.12.021
- Benson, D. & Smith, J. (2015). Protecting biodiversity values in response to long-term impacts: additional areas recommended for inclusion in the Greater Blue Mountains World Heritage Area. In: *Values for a new generation: Greater Blue Mountains World Heritage Area* (ed Benson, D.) pp. 48-75: Greater Blue Mountains World Heritage Area Advisory Committee. Available from <http://www.environment.nsw.gov.au/protectedareas/values-new-generation.htm>.
- Benson, D.H. & Keith, D.A. (1990). Natural vegetation of the Wallerawang 1:100,000 map sheet. *Cunninghamia*, 2, 305-335.
- Binns, D.L. (1997). Floristics and vegetation patterns of Coolah Tops, New South Wales. *Cunninghamia*, 5, 233-274.
- Black, M.P., Mooney, S.D. & Attenbrow, V. (2008). Implications of a 14200 year contiguous fire record for understanding human-climate relationships at Gooches Swamp, New South Wales, Australia. *The Holocene*, 18, 437-447.
- Brown, M.J., Crowden, R.K. & Jarman, S.J. (1982). Vegetation of an alkaline pan-acidic peat mosaic in the Hardwood River Valley, Tasmania. *Austral Ecology*, 7, 3-12.
- Chalson, J.M. & Martin, H.A. (2009). A Holocene history of the vegetation of the Blue Mountains, New South Wales. *Proceedings of the Linnean Society of New South Wales*, 130, 77-109.
- CoA. (2005). Commonwealth listing advice on Temperate Highlands Peat Swamps on Sandstone. [Accessed January 10th 2018]. Retrieved from <http://www.environment.gov.au/biodiversity/threatened/communities/temperate-highland-peat-swamps.html>
- CoA. (2010). Temperate Highland Peat Swamps on Sandstone in Community and Species Profile and Threats Database. [Accessed January 10th 2018]. Retrieved from <http://www.environment.gov.au/sprat>
- Cowardin, L.M., Carter, V., Golet, F.C. & LaRoe, E.T. (1979). *Classification of Wetlands and Deepwater Habitats of the United States*. Washington, D.C.: U.S. Department of the Interior, Fish and Wildlife Service.
- Craven, L.A. (2009). *Melaleuca* (Myrtaceae) from Australia. *Novon*, 19, 444-453.
- DEC. (2006). The Vegetation of the Western Blue Mountains. Unpublished report funded by the Hawkesbury-Nepean Catchment Management Authority. Hurstville: Department of Environment and Conservation NSW.
- DECCW. (2010). *NSW Wetlands Policy*: Department of Environment, Climate Change & Water NSW, Hurstville.
- DLWC. (1999a). Blue Mountains Sandstone Aquifer Current Groundwater Management Practices and Issues. Parramatta, NSW: Department of Land and Water Conservation.
- DLWC. (1999b). Blue Mountains Sandstone Aquifer Status Report. Parramatta, NSW: Department of Land and Water Conservation.

- Douglas, S.M. & Wilson, P.G. (2015). *Callistemon purpurascens*: a new and threatened species from the Blue Mountains region, New South Wales, Australia. *Telopea*, 18, 265-272. doi:10.7751/telopea8562
- Downing, A.J., Brown, E.A., Oldfield, R.J., Selkirk, P.M. & Coveny, R. (2007). Bryophytes and their distribution in the Blue Mountains of New South Wales. *Cunninghamia*, 10, 225-254.
- Ford, A. (1989). The sclerophyllous flora of Wollemi National Park. Unpublished report.
- Fryirs, K., Freidman, B., Williams, R. & Jacobsen, G. (2014). Peatlands in eastern Australia? Sedimentology and age structure of Temperate Highland Peat Swamps on Sandstone (THPSS) in the Southern Highlands and Blue Mountains of NSW, Australia. *Holocene*, 24, 1527-1538. doi:10.1177/0959683614544064
- GBMWAAC. (2015). *Values for a new generation: Greater Blue Mountains World Heritage Area*: Greater Blue Mountains World Heritage Area Advisory Committee. Available from <http://www.environment.nsw.gov.au/protectedareas/values-new-generation.htm>.
- Gellie, N.J.H. & McRae, R.M. (1985). Vegetation of the Western Blue Mountains-Wollemi Region. Unpublished internal report to NSW National Parks and Wildlife Service, Blue Mountains District.
- Gore, A.J.P. (1983). *Ecosystems of the World 4A. Mires: Swamp, Bog, Fen and Moor*. Amsterdam: Elsevier.
- Gorissen, S. (2016). Conservation biology of the endangered Blue Mountains Water Skink (*Eulamprus leuraensis*). PhD thesis. Sydney, Australia: University of Sydney.
- Hájek, M., Horsák, M., Hájková, P. & Dítě, D. (2006). Habitat diversity of central European fens in relation to environmental gradients and an effort to standardise fen terminology in ecological studies. *Perspectives in Plant Ecology, Evolution and Systematics*, 8, 97-114.
- Holland, W.N., Benson, D.H. & McRae, R.H.D. (1992a). Spatial and temporal variation in a perched headwater valley in the Blue Mountains: geology, geomorphology, vegetation, soils and hydrology. *Proceedings of the Linnean Society of New South Wales*, 113, 271-295.
- Holland, W.N., Benson, D.H. & McRae, R.H.D. (1992b). Spatial and temporal variation in a perched headwater valley in the Blue Mountains: solar radiation and temperature. *Proceedings of the Linnean Society of New South Wales*, 113, 297-309.
- Hope, G., Nanson, R. & Flett, I. (2009). Technical Report 19. The peat-forming mires of the Australian Capital Territory. Canberra: Territory and Municipal Services.
- Hunter, J.T. (2013). Upland wetlands in the Namoi Catchment: mapping distribution and disturbance classes of fens, bogs and lagoons. *Cunninghamia*, 13, 331-335.
- Hunter, J.T. & Bell, D. (2007). Vegetation of montane bogs in east-flowing catchments of northern New England, New South Wales. *Cunninghamia*, 10, 77-92.
- Hunter, J.T. & Bell, D. (2009). The *Carex* fen vegetation of northern New South Wales. *Cunninghamia*, 11, 49-64.
- Joosten, H., Tanneberger, F. & Moen, A. (2017). Mires and Peatlands of Europe: Status, distribution and conservation. Stuttgart: Schweizerbart Science Publishers.
- Keith, D.A. (2004). *Ocean Shores to Desert Dunes: the Native Vegetation of New South Wales and the ACT*. Hurstville: Department of Environment and Conservation (NSW).
- Keith, D.A. & Benson, D.H. (1988). Natural vegetation of the Katoomba 1:100,000 map sheet. *Cunninghamia*, 2, 107-143.
- Keith, D.A., Elith, J. & Simpson, C.C. (2014). Predicting distribution changes of a mire ecosystem under future climates. *Diversity and Distributions*, 20, 440-454. doi:10.1111/ddi.12173
- Keith, D.A. & Myerscough, P.J. (1993). Floristics and soil relations of upland swamp vegetation near Sydney. *Australian Journal of Ecology*, 18, 325-344.
- Keith, D.A., Rodoreda, S., Holman, L. & Lemmon, J. (2006). Monitoring change in upland swamps in Sydney's water catchments: the roles of fire and rain. Sydney Catchment Authority Special Area Strategic Management Research and Data Program. Project number RD07: Long term responses of upland swamps to fire. Final Report. Hurstville, NSW: Department of Environment and Conservation.
- Kirkpatrick, J.B. & Bridle, K.L. (1998). Environmental relationships of floristic variation in the alpine vegetation of southeastern Australia. *Journal of Vegetation Science*, 9, 251-260.
- Kodala, P., James, T.A. & Hind, P.D. (1996). Vegetation and flora of swamps of the Boyd Plateau, Central Tablelands, New South Wales. *Cunninghamia*, 4, 525-530.
- Kovac, M. & Lawrie, J.W. (1991). *Soil Landscapes of the Singleton 1:250,000 Sheet*. Sydney: Soil Conservation Service of NSW.
- LeBreton, M. (1996). Habitat and distribution of the Blue Mountains swamp skink (*Eulamprus leuraensis*). B. Zool. (Honours) thesis, University of New South Wales.
- Mactaggart, B., Bauer, J., Goldney, D. & Rawson, A. (2008). Problems in naming and defining the swampy meadow: an Australian perspective. *Journal of Environmental Management*, 87, 461-473.
- Mactaggart, B.G. (2008). Characterising and understanding swampy meadows in the NSW Central Tablelands region: a prerequisite for their restoration. PhD thesis. The University of Sydney.
- Marshall, B. (2005). *Groundwater: Lifeblood of the Environment*. Wentworth Falls, NSW: Blue Mountains Conservation Society.
- Martin, L. (2017). Records of postglacial hydroclimatic variability from the peat-forming environments of the Sydney Region. Unpublished PhD Thesis, School of Biological, Earth and Environmental Sciences. Sydney: University of New South Wales.
- McCormack, R.B. (2012). *A Guide to Australia's Spiny Freshwater Crayfish*. Collingwood, Victoria: CSIRO Publishing.
- McCormack, R.B. & Ahyong, S.T. (2017). *Euastacus vesper* sp. nov., a new giant spiny crayfish (Crustacea, Decapoda, Parastacidae) from the Great Dividing Range, New South Wales, Australia. *Zootaxa*, 4244, 556-567. doi:10.11646/zootaxa.4244.4.6
- Mitsch, W.J. & Gosselink, J.G. (2007). *Wetlands*. Hoboken, New Jersey: John Wiley & Sons.
- Mitsch, W.J., Gosselink, J.G., Anderson, C.J. & Zhang, L. (2009). *Wetland Ecosystems*. Hoboken, New Jersey: John Wiley & Sons.
- Mort, S.J. (1983). The Barrington Tops swamps: flora, ecology and conservation. Unpublished B.Sc. (Hons) thesis. University of New South Wales.
- NSW Government. (2002). *The NSW State Groundwater Dependent Ecosystems Policy*. Sydney: Department of Land and Water Conservation and the State Groundwater Policy Working Group.
- NSWSC. (2004). Montane Peatlands and Swamps of the New England Tableland, NSW North Coast, Sydney Basin, South East Corner, South Eastern Highlands and Australian Alps bioregions - endangered ecological community listing. NSW Scientific Committee final determination. [Accessed January 10th 2018]. Retrieved from <http://www.environment.nsw.gov.au/determinations/MontanePeatlandsEndSpListing.htm>
- NSWSC. (2005a). Alteration of habitat following subsidence due to longwall mining. NSW Scientific Committee Key Threatening Process final determination. [Accessed January 10th 2018]. Retrieved from <http://www.environment.nsw.gov.au/determinations/LongwallMiningKtp.htm>

- NSWSC. (2005b). Newnes Plateau Shrub Swamp in the Sydney Basin Bioregion - endangered ecological community listing. NSW Scientific Committee final determination. [Accessed January 10th 2018]. Retrieved from <http://www.environment.nsw.gov.au/determinations/NewnesPlateauShrubSwampEndSpListing.htm>
- NSWSC. (2007). Blue Mountains Swamps in the Sydney Basin Bioregion - vulnerable ecological community listing. NSW Scientific Committee final determination. [Accessed January 10th 2018]. Retrieved from <http://www.environment.nsw.gov.au/determinations/BlueMountainsSwampsVulnerableEcologicalCommunity.htm>
- Pemberton, M. (2005). Australian peatlands: a brief consideration of their origin, distribution, natural values and threats. *Journal of the Royal Society of Western Australia*, 88, 81-89.
- Rydin, H. & Jeglum, J.K. (2013). *The Biology of Peatlands*. Oxford: Oxford University Press.
- Serov, P., Kuginis, L. & Williams, J.P. (2012). *Risk assessment guidelines for groundwater dependent ecosystems, Volume 1 - the conceptual framework*. Sydney: Department of Primary Industries, NSW Office of Water.
- Shepherd, J. & Keyzer, V. (2014). Ecology of *Eucalyptus aquatica* (Myrtaceae), a restricted eucalypt confined to montane swamp (fen) habitat in south-eastern Australia. *Cunninghamia*, 14, 63-76. doi:10.7751/cunninghamia.2014.14.004
- Tame, T. (1997). The vegetation of the Dunns Swamp Area. Unpublished report to NSW National Parks and Wildlife Service, Mudgee Sub-district.
- Thackway, R. & Cresswell, I.D. (1995). *An Interim Biogeographic Regionalisation for Australia: a Framework for Establishing the National System of Reserves*. Canberra: Australian Nature Conservation Agency.
- Tozer, M.G., Turner, K., Keith, D.A., Tindall, D., Pennay, C., Simpson, C. & Mackenzie, B. (2010). Native vegetation of southeast NSW: a revised classification and map for the coast and eastern tablelands. *Cunninghamia*, 11, 359-406.
- Udovicic, F. & Spencer, R.D. (2012). New combinations of *Callistemon* (Myrtaceae). *Muelleria*, 30, 23-25.
- Wheeler, B.D. & Proctor, M.C.F. (2000). Ecological gradients, subdivisions and terminology of north-west European mires. *Journal of Ecology*, 88, 187-203. doi:10.1046/j.1365-2745.2000.00455.x
- Whinam, J. (1995). Effects of fire on Tasmanian *Sphagnum* peatlands. In: *Bushfire '95: Presented papers, Australian Bushfire Conference, 27-30 September 1995* pp. 1-13. Hobart, Tasmania: Forestry Tasmania, Parks and Wildlife Service, Tasmanian Fire Service.
- Whinam, J. & Chilcott, N. (2002). Floristic description and environmental relationships of *Sphagnum* communities in NSW and the ACT and their conservation management. *Cunninghamia*, 7, 463-500.
- Whinam, J., Eberhard, S., Kirkpatrick, J. & Moscal, T. (1989). *Ecology and Conservation of Tasmanian Sphagnum Peatlands*. Hobart: Tasmanian Conservation Trust.
- Whinam, J. & Hope, G.S. (2005). The peatlands of the Australasian region. In: *Moore: von Sibirien bis Feuerland - Mires: from Siberia to Tierra del Fuego* (ed Steiner, G.M.) pp. 397-434. Linz, Austria: Biologiezentrum der Oberoesterreichischen Landesmuseen Neue Serie 35.
- Yabe, K. & Onimaru, K. (1997). Key variables controlling the vegetation of a cool-temperate mire in northern Japan. *Journal of Vegetation Science*, 8, 29-36.

Appendix 1: Native plant species (including mosses) recorded in Rollen Creek mire (TB=Tableland Bog, TSM=Tableland Swamp Meadow, HS= Hanging Swamp), and whether those species have also been recorded in Newnes Plateau Shrub Swamps (NPSS) (Benson & Baird, 2012) or Boyd Plateau Bogs (Kodela et al., 1996).

PLANT SPECIES	Family	Rollen Creek mire	NPSS	Boyd Plateau Bogs
<i>Asperula gunnii</i>	Rubiaceae	TB	NPSS	BPB
<i>Baeckea utilis</i>	Myrtaceae	TB, HS	NPSS	BPB
<i>Baloskion australe</i>	Restionaceae	TB, HS	NPSS	BPB
<i>Baumea</i> sp.	Cyperaceae	TB, HS	NPSS	BPB
<i>Blechnum nudum</i>	Blechnaceae	TB	NPSS	BPB
<i>Bulbine bulbosa</i>	Liliaceae	TB		
<i>Callistemon citrinus</i>	Myrtaceae	TB, HS		
<i>Carex gaudichaudiana</i>	Cyperaceae	TB, TSM		BPB
<i>Centella asiatica</i> (or <i>cordifolia</i>)	Apiaceae	TB		BPB
<i>Centrolepis?</i>	Centrolepiaceae	TB		
<i>Comesperma retusum</i>	Polygalaceae	TB		BPB
<i>Drosera binata</i>	Droseraceae	TB, HS	NPSS	BPB
<i>Drosera spathulata</i>	Droseraceae	TB	NPSS	
<i>Empodisma minus</i>	Restionaceae	TB, HS	NPSS	BPB
<i>Epacris microphylla</i>	Ericaceae	TB	NPSS	BPB
<i>Epacris paludosa</i>	Ericaceae	TB, HS	NPSS	BPB
<i>Epilobium gunnianum</i>	Onagraceae			BPB
<i>Eriocaulon scariosum</i>	Eriocaulaceae	TB		
<i>Eucalyptus camphora</i> subsp. <i>camphora</i>	Myrtaceae	TB, HS		
<i>Eucalyptus pauciflora</i>	Myrtaceae	TB	NPSS	BPB
<i>Gahnia sieberiana</i>	Cyperaceae		NPSS	
<i>Geranium neglectum</i>	Geraniaceae	TB		BPB
<i>Gleichenia dicarpa</i>	Gleicheniaceae	TB, HS	NPSS	BPB
<i>Gonocarpus micrantha</i>	Haloragaceae	TB	NPSS	BPB
<i>Goodenia</i> sp. (<i>bellidifolia</i>)	Goodeniaceae	TB	NPSS	
<i>Gymnoschoenus sphaerocephalus</i>	Cyperaceae	TB, HS	NPSS	
<i>Hakea microcarpa</i>	Proteaceae	TB	NPSS	BPB
<i>Hybanthus?</i>	Violaceae	TB		
<i>Hydrocotyle</i> sp. (<i>sibthorpioides?</i>)	Apiaceae	TB	NPSS	BPB
<i>Hypericum</i> sp. (<i>gramineum?</i>)	Hypericaceae	TB	NPSS	
<i>Isachne globosa</i>	Poaceae	TB		
<i>Isotoma fluviatilis?</i> = <i>Pratia surrepens</i>	Campanulaceae	TB		BPB
<i>Juncus</i> spp.	Juncaceae	TB, TSM	NPSS	BPB
<i>Lepidosperma limicola</i>	Cyperaceae	TB, HS	NPSS	
<i>Leptospermum continentale</i>	Myrtaceae	TB	NPSS	BPB
<i>Leptospermum myrtifolium</i>	Myrtaceae		NPSS	BPB
<i>Leptospermum obovatum</i>	Myrtaceae	TB, HS	NPSS	BPB
<i>Lepyrodia</i> spp.	Restionaceae	TB	NPSS	
<i>Patersonia fragilis</i>	Iridaceae	TB	NPSS	BPB
<i>Pultenea divaricata</i>	Fabaceae	TB, HS	NPSS	
<i>Pultenea</i> sp.	Fabaceae	TB		
<i>Ranunculus</i> sp.	Ranunculaceae	TB		
<i>Scirpus polystachyus</i>	Cyperaceae			BPB
<i>Senecio</i> sp.	Asteraceae	TB		
<i>Sphagnum cristatum</i>	Sphagnaceae	TB, HS		BPB
<i>Spiranthes australis</i>	Orchidaceae	TB		BPB
<i>Stylidium graminifolium</i>	Stylidiaceae	TB	NPSS	BPB
<i>Tetrarrhena juncea</i>	Poaceae	TB	NPSS	
<i>Utricularia dichotoma</i>	Lentibulariaceae	TB	NPSS	BPB
<i>Viola caleyana</i>	Violaceae	TB	NPSS	
<i>Xyris ustulata</i>	Xyridaceae	TB, HS	NPSS	BPB
<i>Xyris gracilis</i>	Xyridaceae	TB	NPSS	
TOTAL		48	33	31
Number RC shared with NPSS and/or BPB			31	28
Percentage RC shared with NPSS and/or BPB			65%	58%

